

TERNARY UNIVERSAL SUMS OF GENERALIZED POLYGONAL NUMBERS

JANGWON JU, BYEONG-KWEON OH AND BANGNAM SEO

ABSTRACT. An integer of the form $p_m(x) = \frac{(m-2)x^2 - (m-4)x}{2}$ ($m \geq 3$), for some integer x is called a generalized polygonal number of order m . A ternary sum $\Phi_{i,j,k}^{a,b,c}(x, y, z) = ap_i(x) + bp_j(y) + cp_k(z)$ of generalized polygonal numbers, for some positive integers a, b, c and some integers $i, j, k \geq 3$, is said to be *universal* if for any positive integer n , the equation $\Phi_{i,j,k}^{a,b,c}(x, y, z) = n$ has an integer solution x, y, z . In this article, we prove the universalities of 17 ternary sums of generalized polygonal numbers listed in (1.2). This was conjectured by Sun in [11].

1. INTRODUCTION

A polygonal number is a number represented as dots or pebbles arranged in the shape of a regular polygon. More precisely, a *polygonal number of order m* (or an *m -gonal number*) for $m \geq 3$ is defined by

$$p_m(x) = \frac{(m-2)x^2 - (m-4)x}{2}$$

for some non negative integer x . If we admit that x is a negative integer, then $p_m(x)$ is called a *generalized polygonal number of order m* (or a *generalized m -gonal number*). By definition, every polygonal number of order m is a generalized polygonal number of order m . However, the converse is true only when $m = 3, 4$.

A famous assertion of Fermat states that every positive integer can be expressed as a sum of three triangular numbers, which was proved by Gauss in 1796. This follows from the Gauss-Legendre theorem, which states that every positive integer which is not of the form $4^k(8l+7)$ with non negative integers k and l , is a sum of three squares of integers.

In general, a ternary sum $\Phi_{i,j,k}^{a,b,c} = ap_i(x) + bp_j(y) + cp_k(z)$ ($a, b, c > 0$) of polygonal numbers is called *universal over $\mathbb{N}$* if for any non negative integer n , the diophantine equation

$$(1.1) \quad ap_i(x) + bp_j(y) + cp_k(z) = n$$

2000 *Mathematics Subject Classification.* Primary 11E12, 11E20.

Key words and phrases. generalized polygonal numbers, ternary universal sums.

This work of the first author was supported by BK21 PLUS SNU Mathematical Sciences Division.

This work of the second author was supported by the National Research Foundation of Korea (NRF-2014R1A1A2056296).

has a *non negative* integer solution x, y, z . It is called *universal over $\mathbb{Z}$* if the above equation has an integer solution.

In 1862, Liouville generalized Gauss's triangular theorem as follows: for positive integers a, b, c ($a \leq b \leq c$), the ternary sum $ap_3(x) + bp_3(y) + cp_3(z)$ of triangular numbers is universal if and only if (a, b, c) is one of the following triples:

$$(1, 1, 1), (1, 1, 2), (1, 1, 4), (1, 1, 5), (1, 2, 2), (1, 2, 3), (1, 2, 4).$$

In [3], [8] and [10], Sun and his collaborators determined all ternary universal sums of the form $ap_3(x) + bp_3(y) + cp_4(z)$ or $ap_3(x) + bp_4(y) + cp_4(z)$. There does not exist a universal sum of the form $ap_4(x) + bp_4(y) + cp_4(z)$ for any positive integers a, b, c . Hence, to find all ternary sums of polygonal numbers that are universal over $\mathbb{N}$, we may assume that $\{i, j, k\} \neq \{3, 4\}$. Recently, Sun gave in [11] a complete list of 95 candidates of ternary sums of polygonal numbers that are universal over $\mathbb{N}$ under this assumption.

In fact, the proof of the universality over $\mathbb{Z}$ of a ternary sum of polygonal numbers is much easier than the proof of universality over $\mathbb{N}$. For this reason, one may naturally ask the universalities over $\mathbb{Z}$ of 95 candidates of ternary sums. Related with this question, Sun and Ge proved in [2] and [11] that 62 ternary sums among 95 candidates are, in fact, universal over $\mathbb{Z}$. The remaining $33 = 95 - 62$ candidates are as follows:

$$(1.2) \quad \begin{aligned} & \mathbf{p_3} + \mathbf{9p_3} + \mathbf{p_5}, \mathbf{p_3} + \mathbf{2p_4} + \mathbf{4p_5}, \mathbf{p_3} + \mathbf{4p_4} + \mathbf{2p_5}, \mathbf{p_3} + \mathbf{2p_3} + \mathbf{p_7}, \\ & \mathbf{p_3} + \mathbf{p_4} + \mathbf{2p_7}, \mathbf{p_3} + \mathbf{p_5} + \mathbf{p_7}, \quad p_3 + p_5 + 4p_7, \quad \mathbf{p_3} + \mathbf{2p_5} + \mathbf{p_7}, \\ & \mathbf{p_3} + \mathbf{p_7} + \mathbf{2p_7}, \mathbf{p_3} + \mathbf{2p_3} + \mathbf{2p_8}, \mathbf{p_3} + \mathbf{p_7} + \mathbf{p_8}, \quad \mathbf{p_3} + \mathbf{2p_3} + \mathbf{p_9}, \\ & p_3 + 2p_3 + 2p_9, \mathbf{p_3} + \mathbf{p_4} + \mathbf{p_9}, \quad p_3 + p_4 + 2p_9, \quad p_3 + 2p_4 + p_9, \\ & \mathbf{p_3} + \mathbf{p_5} + \mathbf{2p_9}, \mathbf{2p_3} + \mathbf{p_5} + \mathbf{p_9}, \quad \mathbf{p_3} + \mathbf{p_3} + \mathbf{p_{12}}, \quad \mathbf{p_3} + \mathbf{2p_3} + \mathbf{p_{12}}, \\ & p_3 + 2p_3 + 2p_{12}, p_3 + p_4 + p_{13}, \quad \mathbf{p_3} + \mathbf{p_5} + \mathbf{p_{13}}, \quad p_3 + 2p_3 + p_{15}, \\ & p_3 + p_4 + p_{15}, \quad p_3 + 2p_3 + p_{16}, \quad p_3 + p_3 + p_{17}, \quad p_3 + 2p_3 + p_{17}, \\ & p_3 + p_4 + p_{17}, \quad p_3 + 2p_4 + p_{17}, \quad p_3 + p_4 + p_{18}, \quad p_3 + 2p_3 + p_{23}, \\ & p_3 + p_4 + p_{27}. \end{aligned}$$

In this paper, we prove that 17 ternary sums written in boldface in the above remaining candidates are universal over $\mathbb{Z}$. In most cases, we use the method developed in [7]. For those who are unfamiliar with it, we briefly review this method in Section 2.

The term “lattice” will always refer to a positive definite integral $\mathbb{Z}$ -lattice on an n -dimensional positive definite quadratic space over $\mathbb{Q}$. Let $L = \mathbb{Z}x_1 + \cdots + \mathbb{Z}x_n$ be a $\mathbb{Z}$ -lattice of rank n . The corresponding symmetric matrix M_L is defined by

$$M_L = (B(x_i, x_j)).$$

If M_L is diagonal, then we simply write $L = \langle Q(x_1), \dots, Q(x_n) \rangle$. The genus of L is the set of all lattices that are locally isometric to L , which is denoted by $\text{gen}(L)$. The subset of $\text{gen}(L)$ consisting of all lattices that are isometric to L itself is denoted by $[L]$. An integer a is called an *eligible integer of L* if it is locally represented by L . Note that any eligible integer of L is represented a lattice in the genus of L . The set of all eligible integers of L is denoted by $Q(\text{gen}(L))$. Similarly, $Q(L)$ denotes

the set of all integers that are represented by L itself. For an integer a , $R(a, L)$ denotes the set of all vectors $x \in L$ such that $Q(x) = a$, and $r(a, L)$ denotes its cardinality. For a set S , we define $\pm S = \{s : s \in S \text{ or } -s \in S\}$. For two vectors $v = (v_1, v_2, \dots, v_n), w = (w_1, w_2, \dots, w_n) \in \mathbb{Z}^n$ and a positive integer s , if $v_i \equiv w_i \pmod{s}$ for any $i = 1, 2, \dots, n$, then we write $v \equiv w \pmod{s}$.

Any unexplained notations and terminologies can be found in [5] or [9].

2. GENERAL TOOLS

Let i, j and k be integers with $1 \leq i \leq j \leq k$. Recall that for positive integers a, b and c , a ternary sum

$$\Phi_{i,j,k}^{a,b,c}(x, y, z) = ap_{i+2}(x) + bp_{j+2}(y) + cp_{k+2}(z)$$

of generalized polygonal numbers is universal over $\mathbb{Z}$ if $\Phi_{i,j,k}^{a,b,c}(x, y, z) = n$ has an integer solution for any non negative integer n . Since any generalized hexagonal number is a triangular number and vice versa, we always assume that neither i, j nor k is 4. Note that $\Phi_{i,j,k}^{a,b,c}$ is universal over $\mathbb{Z}$ if and only if the equation

$$(2.1) \quad \begin{aligned} & jka(2ix - (i-2))^2 + ikb(2jy - (j-2))^2 + ijc(2kz - (k-2))^2 \\ & = 8ijkn + jka(i-2)^2 + ikb(j-2)^2 + ijc(k-2)^2 =: \Psi_{i,j,k}^{a,b,c}(n) \end{aligned}$$

has an integer solution for any non negative integer n . This is equivalent to the existence of an integer solution X, Y and Z of the diophantine equation

$$(2.2) \quad jkaX^2 + ikbY^2 + ijcZ^2 = \Psi_{i,j,k}^{a,b,c}(n)$$

satisfying the following congruence condition

$$(2.3) \quad X \equiv -i + 2 \pmod{2i}, Y \equiv -j + 2 \pmod{2j} \text{ and } Z \equiv -k + 2 \pmod{2k}.$$

In some particular cases, representations of quadratic forms with some congruence conditions corresponds to representations of a subform which is taken suitably (for details, see [4]). The following lemma reveals this correspondence more concretely in our case.

Lemma 2.1. *Assume that i, j, k and a, b, c given above satisfy the followings:*

- (i) $\{i, j, k\} \subset \mathcal{S}_P = \{1, 2, p^n, 2p^n : p \text{ is an odd prime and } n \geq 1\}$;
- (ii) all of $(i, jka), (j, ikb)$ and (k, ijc) are one or two;
- (iii) either i, j, k are all even or some of them are odd and

$$\Psi_{i,j,k}^{a,b,c}(n) \not\equiv 0 \pmod{2^{3+\text{ord}_2(ijk)}}.$$

Then there is a quadratic form $Q(x, y, z)$ such that for any integer n , $Q(x, y, z) = \Psi_{i,j,k}^{a,b,c}(n)$ has an integer solution if and only if Equation (2.2) satisfying (2.3) has an integer solution.

Proof. Note that for any $s \in \mathcal{S}_P$ and any integer a with $(a, s) = 1$, the congruence equation $x^2 \equiv a^2 \pmod{s}$ has at most two incongruent integer solutions modulo s .

First, assume that all of i, j and k are even. From the assumption, we have $i \equiv j \equiv k \equiv 2 \pmod{4}$. Define

$$Q(x, y, z) = 16(jkaX^2 + ikbY^2 + ijcZ^2).$$

Assume that $(u, v, w) \in \mathbb{Z}^3$ is an integer solution of $Q(x, y, z) = \Psi_{i,j,k}^{a,b,c}(n)$. Then

$$16jka u^2 \equiv jka(i-2)^2 \pmod{32i}.$$

From this and the conditions (i) and (ii) follows $4u \equiv \pm(i-2) \pmod{2i}$. Similarly, we have $4v \equiv \pm(j-2) \pmod{2j}$ and $4k \equiv \pm(k-2) \pmod{2k}$. Therefore by choosing the signs suitably, we may find an integer solution of Equation (2.2) satisfying (2.3). For the converse, note that any integers X, Y, Z satisfying the condition (2.3) are divisible by 4.

Assume that exactly one of i, j, k , say i , is odd. Define

$$Q(x, y, z) = jkaX^2 + 16(ikbY^2 + ijcZ^2).$$

Assume that $(u, v, w) \in \mathbb{Z}^3$ is an integer solution of $Q(x, y, z) = \Psi_{i,j,k}^{a,b,c}(n)$. Since

$$16ikbv^2 \equiv ikb(j-2)^2 \pmod{j} \quad \text{and} \quad 16ijcw^2 \equiv ijc(k-2)^2 \pmod{k},$$

and $(j, ikb) = (k, ijc) = 2$ from (ii), we have

$$16v^2 \equiv (j-2)^2 \pmod{\frac{j}{2}} \quad \text{and} \quad 16w^2 \equiv (k-2)^2 \pmod{\frac{k}{2}},$$

which implies that $4v \equiv \pm(j-2) \pmod{2j}$ and $4k \equiv \pm(k-2) \pmod{2k}$. Now since $jka u^2 \equiv jka(i-2)^2 \pmod{32i}$ and $a \not\equiv 0 \pmod{8}$ by the condition (iii), we have $u^2 \equiv (i-2)^2 \pmod{2i}$. Hence $u \equiv \pm(i-2) \pmod{2i}$. Therefore by choosing the signs suitably, we may find an integer solution of (2.2) satisfying (2.3).

Assume that exactly two of i, j, k , say j, k , are odd. Define

$$Q(x, y, z) = 16jkaX^2 + ikbY^2 + ijc(Y-4Z)^2.$$

Assume that $(u, v, w) \in \mathbb{Z}^3$ is an integer solution of $Q(x, y, z) = \Psi_{i,j,k}^{a,b,c}(n)$. Then, similarly to the above, we may easily show that

$$4u \equiv \pm(i-2) \pmod{\frac{i}{2}}, \quad v \equiv \pm(j-2) \pmod{j}, \quad v-4w \equiv \pm(k-2) \pmod{k}.$$

Since $i \equiv 2 \pmod{4}$, we have $4u \equiv \pm(i-2) \pmod{2i}$. Furthermore, since

$$ikbv^2 + ijc(v-4w)^2 \equiv ikb + ijc \not\equiv 0 \pmod{16},$$

v is odd. Hence

$$v \equiv \pm(j-2) \pmod{2j}, \quad v-4w \equiv \pm(k-2) \pmod{2k}.$$

Therefore by choosing the signs suitably, we may find an integer solution of (2.2) satisfying (2.3). Conversely, if Equation (2.2) satisfying (2.3) has an integer solution X, Y, Z , then X is divisible by 4 and both Y and Z are odd. Hence $Y \equiv Z \pmod{4}$ or $Y \equiv -Z \pmod{4}$. This implies that $Q(x, y, z) = \Psi_{i,j,k}^{a,b,c}(n)$ has an integer solution.

Finally, if all of i, j, k are odd, then we define

$$Q(x, y, z) = jkaX^2 + ikb(X-4Y)^2 + ijc(X-4Z)^2.$$

The proof of the remaining part is quite similar to the above. $\square$

Remark 2.2. Among 33 candidates given in the introduction, those who do not satisfy the conditions in Lemma 2.1 are the cases

$$4, 7, 9, 14, 16, 24, \text{ and } 27 \sim 32.$$

For 4-th, 9-th and 14-th cases among them, we prove their universalities by using some other methods. For details, see Section 3.

For the representation of an arithmetic progression by a ternary quadratic form, we use the method developed in [6] and [7]. For those who are unfamiliar with this, we briefly introduce how it works.

Let d be a positive integer and let a be a non negative integer ($a \leq d$). We define

$$S_{d,a} = \{dn + a \mid n \in \mathbb{N} \cup \{0\}\}.$$

For ternary $\mathbb{Z}$ -lattices M, N on the quadratic space V , we define

$$R(N, d, a) = \{x \in N/dN \mid Q(x) \equiv a \pmod{d}\}$$

and

$$R(M, N, d) = \{\sigma \in O(V) \mid \sigma(dN) \subset M\}.$$

A coset (or, a vector in the coset) $v \in R(N, d, a)$ is said to be *good* with respect to M, N, d and a if for any $x \in N$ satisfying $x \equiv v \pmod{d}$, there is a $\sigma \in R(M, N, d)$ such that $\sigma(x) \in M$. If every vector contained in $R(N, d, a)$ is good, we write

$$N <_{d,a} M.$$

The set of all good vectors in $R(N, d, a)$ is denoted by $R_M(N, d, a)$. If $N <_{d,a} M$, then by Lemma 2.2 of [6], we have

$$S_{d,a} \cap Q(N) \subset Q(M).$$

Note that the converse is not true in general.

The following theorem is a slightly modified version of Corollary 2.2 of [7]:

Theorem 2.3. *Under the same notations given above, assume that there is a partition $R(N, d, a) - R_M(N, d, a) = (P_1 \cup \cdots \cup P_k) \cup (\tilde{P}_1 \cup \cdots \cup \tilde{P}_{k'})$ satisfying the following properties: for each P_i , there is a $\tau_i \in O(V)$ such that*

- (i) τ_i has an infinite order;
- (ii) $\tau_i(dN) \subset N$;
- (iii) for any $v \in P_i$, $\tau_i(v) \in N$ and $\tau_i(v) \pmod{d} \in P_i \cup R_M(N, d, a)$,

and for each $\tilde{P}_j$, there is a $\tilde{\tau}_j \in O(V)$ such that

- (iv) $\tilde{\tau}_j(dN) \subset N$;
- (v) for any $v \in \tilde{P}_j$, $\tilde{\tau}_j(v) \in N$ and $\tilde{\tau}_j(v) \pmod{d} \in P_1 \cup \cdots \cup P_k \cup R_M(N, d, a)$.

Then we have

$$(S_{d,a} \cap Q(N)) - \bigcup_{i=1}^k Q(z_i)\mathbb{Z}^2 \subset Q(M),$$

where the vector z_i is a primitive vector in N which is an eigenvector of τ_i .

Proof. From the conditions (iv) and (v), we may assume that

$$R(N, d, a) - R_M(N, d, a) = P_1 \cup \cdots \cup P_k.$$

Hence the theorem follows directly from Corollary 2.2 in [7]. $\square$

Lemma 2.4. *Let L be a ternary $\mathbb{Z}$ -lattice with class number greater than 1. Assume that the genus of L contains only one spinor genus. If an integer a is represented by all isometric classes in the genus of L except at most only one isometric class, then for any positive integer $s > 1$ with $(s, dL) = 1$, as^2 is represented by all lattices in the genus of L .*

Proof. Without loss of generality, we may assume that there is a $\mathbb{Z}$ -lattice, say M , that does not represent a . For any prime p not dividing dL , the graph $Z(p, L)$ defined in [1] is connected. Hence there is a $\mathbb{Z}$ -lattice M' such that M' is adjacent to M and a is represented by M' . Since pM' is represented by M , ap^2 is represented by M . This completes the proof. $\square$

3. TERNARY UNIVERSAL SUMS OF POLYGONAL NUMBERS

In this section, we prove that 17 ternary sums of generalized polygonal numbers among 33 candidates are, in fact, universal over $\mathbb{Z}$. For the list of all candidates, see (1.2) in the introduction.

(i) $p_3(x) + 9p_3(y) + p_5(z)$. First, we define

$$M = \langle 4, 12, 27 \rangle, \quad N = \langle 3, 4, 12 \rangle.$$

Note that the class number of N is one and every integer of the form $24n + 7$ is represented by N . One may easily show by a direct computation that $(v_1, v_2, v_3) \in R(N, 24, 7) - R_M(N, 24, 7)$ if and only if $v_1, v_2 \equiv \pm 1 \pmod{6}$, $v_3 \equiv 0 \pmod{6}$ and satisfies the following properties:

$$\begin{cases} v_1 \equiv v_2 + v_3 \pmod{4} & \text{if } v_1 \equiv v_2 \pmod{6}, \\ v_1 + v_2 + v_3 \equiv 0 \pmod{4} & \text{otherwise.} \end{cases}$$

Note that $R(N, 24, 7) - R_M(N, 24, 7)$ contains exactly 128 vectors. Define

$$R(N, 24, 7) - R_M(N, 24, 7) = P_1 \cup \tilde{P}_1,$$

where

$$P_1 = \{(v_1, v_2, v_3) \in R(N, 24, 7) - R_M(N, 24, 7) \mid v_1 + v_2 + v_3 \equiv 0 \pmod{4}\},$$

$$\tilde{P}_1 = \{(v_1, v_2, v_3) \in R(N, 24, 7) - R_M(N, 24, 7) \mid v_1 \equiv v_2 + v_3 \pmod{4}\},$$

and

$$\tau_1 = \frac{1}{24} \begin{pmatrix} 0 & 24 & 24 \\ 18 & -6 & 18 \\ -6 & -6 & 18 \end{pmatrix}, \quad \tilde{\tau}_1 = \frac{1}{24} \begin{pmatrix} 0 & -24 & -24 \\ 18 & 6 & -18 \\ -6 & 6 & -18 \end{pmatrix}.$$

Then it satisfies all conditions in Theorem 2.3. The eigenvector of τ_1 is $z_1 = (1, -1, 0)$ and $\tilde{\tau}_1(-1, -1, 0) = (1, -1, 0)$. Hence if $24n + 7$ is not of the form $Q(\pm t, \mp t, 0) = 7t^2$ for some positive integer t , then it is also represented by M .

Assume that $24n + 7 = 7t^2$ for a positive integer t and there is a prime $p > 3$ dividing t . Since the class number of N is one, we have

$$r(7p^2, N) = 4p + 4 - 4 \cdot \left(\frac{-7}{p} \right) > 4$$

by the Minkowski-Siegel formula. Hence there is a representation which is neither contained in the eigenspace of τ_1 nor the inverse image of the eigenspace by $\tilde{\tau}_1$. Therefore every integer of the form $24n + 7$ for a positive integer n is represented by M , that is, the equation

$$24n + 31 = (Y - 4X)^2 + 3Y^2 + 27Z^2 = 4(Y - X)^2 + 12X^2 + 27Z^2$$

has an integer solution X, Y, Z for any integer n . Since $Y - 4X \equiv Y \equiv Z \equiv 1 \pmod{2}$ and $Y - 4X \equiv \pm 1 \pmod{3}$, the equation

$$3(2x + 1)^2 + 27(2y + 1)^2 + (6z - 1)^2 = 24n + 31$$

has an integer solution for any non negative integer n , which implies that the ternary sum $p_3(x) + 9p_3(y) + p_5(z)$ is universal over $\mathbb{Z}$ by (2.1).

(ii) $p_3(x) + 2p_4(y) + 4p_5(z)$. Let

$$M = \langle 3, 4, 48 \rangle, \quad N = \langle 3, 4, 12 \rangle.$$

Since $S_{24,7} \subset Q(\text{gen}(N))$ and $h(N) = 1$, we have $S_{24,7} \subset Q(N)$. One may easily show that

$$R(N, 4, 3) - R_M(N, 4, 3) = \pm\{(1, 0, 1), (1, 0, 3), (1, 2, 1), (1, 2, 3)\}.$$

Let $P_1 = \pm\{(1, 0, 1), (1, 2, 3)\}$, $\tilde{P}_1 = \pm\{(1, 0, 3), (1, 2, 1)\}$ and

$$\tau_1 = \frac{1}{4} \begin{pmatrix} 0 & -4 & -4 \\ 3 & 1 & -3 \\ -1 & 1 & -3 \end{pmatrix}, \quad \tilde{\tau}_1 = \frac{1}{4} \begin{pmatrix} 0 & 4 & 4 \\ 3 & -1 & 3 \\ -1 & -1 & 3 \end{pmatrix}.$$

Then one may easily show that it satisfies all conditions in Theorem 2.3. The eigenvector of τ_1 is $z_1 = (1, 0, 1)$ and $Q(z_1) = 15$, which is not of the form $24n + 7$. Therefore we have $S_{24,7} \subset Q(M)$. This implies that the equation $3X^2 + 4Y^2 + 48Z^2 = 24n + 7$ has an integer solution X, Y, Z for any non negative integer n . Since X is odd and Y is relatively prime to 6, the equation

$$3(2x + 1)^2 + 48y^2 + 4(6z - 1)^2 = 24n + 7$$

has an integer solution for any non negative integer n . Hence by (2.1), the ternary sum $p_3(x) + 2p_4(y) + 4p_5(z)$ is universal over $\mathbb{Z}$.

(iii) $p_3(x) + 4p_4(y) + 2p_5(z)$. Let

$$M = \langle 2, 3, 96 \rangle, \quad N = \begin{pmatrix} 5 & 1 & 1 \\ 1 & 11 & -1 \\ 1 & -1 & 11 \end{pmatrix} \quad \text{and} \quad N' = \langle 2 \rangle \perp \begin{pmatrix} 12 & 6 \\ 6 & 27 \end{pmatrix}.$$

Note that $\text{gen}(M) = \{[M], [N], [N']\}$ and $S_{24,5} \subset Q(\text{gen}(M))$. One may easily show that $N' <_{4,1} N$. Furthermore, $R(N, 12, 5) - R_M(N, 12, 5)$ consists of following vectors:

$$\begin{aligned} & \pm(3, 1, 7), \quad \pm(3, 2, 8), \quad \pm(3, 4, 10), \quad \pm(3, 5, 11), \\ & \pm(3, 7, 1), \quad \pm(3, 8, 2), \quad \pm(3, 10, 4), \quad \pm(3, 11, 5). \end{aligned}$$

If we define $P_1 = R(N, 12, 5) - R_M(N, 12, 5)$ and $\tau_1 = \frac{1}{12} \begin{pmatrix} 8 & 10 & -10 \\ -6 & -3 & -9 \\ 2 & -11 & -1 \end{pmatrix}$,

then P_1 and τ_1 satisfies all conditions in Theorem 2.3. Since the eigenvector of τ_1 is $z_1 = (0, 1, 1)$ and $Q(z_1) = 20$, which is not of the form $24n + 5$, we have $S_{24,5} \cap Q(N) \subset Q(M)$. Therefore every integer of the form $24n + 5$, for any non negative integer n is represented by M . If the equation $2X^2 + 3Y^2 + 96Z^2 = 24n + 5$ has an integer solution X, Y, Z , then X is relatively prime to 6 and Y is odd. Hence the equation

$$3(2x - 1)^2 + 96y^2 + 2(6z - 1)^2 = 24n + 5$$

has an integer solution for any non negative integer n . Therefore the ternary sum $p_3(x) + 4p_4(y) + 2p_5(z)$ is universal over $\mathbb{Z}$.

(iv) $p_3(x) + 2p_3(y) + p_7(z)$. First, we show that for any eligible integer $8n$ of $F(X, Y, Z) = X^2 + 5Y^2 + 10Z^2$, the equation $F(X, Y, Z) = 8n$ has an integral solution $(X, Y, Z) = (a, b, c)$ with $a \equiv b \equiv c \equiv 1 \pmod{2}$, unless $n \equiv 0 \pmod{5}$. Since the class number of F is 1, $F(X, Y, Z) = 8n$ has an integer solution $(X, Y, Z) = (a, b, c)$. One may easily check that all of a, b and c have same parity. Suppose that a, b and c are all even. Then $a \equiv b \pmod{4}$ and there is another representation

$$F\left(\frac{a + 5b + 10c}{4}, \frac{-a + 3b - 2c}{4}, \frac{-a - b + 2c}{4}\right) = F(a, b, c) = 8n.$$

Since there are only finitely many representations of $8n$ and $\tau = \frac{1}{4} \begin{pmatrix} 1 & 5 & 10 \\ -1 & 3 & -2 \\ -1 & -1 & 2 \end{pmatrix}$

has an infinite order, there is a positive integer m such that each component of the vector $\tau^m(a, b, c) = (a_m, b_m, c_m)$ is odd, unless (a, b, c) is not an eigenvector of τ_1 . Note that $(a, b, c) = (0, -2t, t)$ is an eigenvector of τ and $F(0, -2t, t) = 30t^2$.

Note that any positive integer of the form $40n + 24$ for some non negative integer n is an eligible integer of F and the class number of F is one. Hence the equation $F(X, Y, Z) = 40n + 24$ has an integer solution X, Y, Z such that $X \equiv Y \equiv Z \equiv 1 \pmod{2}$ and $X \equiv \pm 2 \pmod{5}$, which implies that

$$5(2x - 1)^2 + 10(2y - 1)^2 + (10z - 3)^2 = 40n + 24$$

has an integer solution for any non negative integer n . Therefore by (2.1), the ternary sum $p_3(x) + 2p_3(y) + p_7(z)$ is universal over $\mathbb{Z}$.

(v) $p_3(x) + p_4 + 2p_7(z)$. Let

$$M = \langle 2, 5, 40 \rangle, \quad N = \langle 5, 8, 10 \rangle.$$

Note that $\text{gen}(M) = \{[M], [N]\}$. The set $R(N, 8, 7) - R_M(N, 8, 7)$ consists of following vectors:

$$\begin{aligned} & \pm(1, 0, 1), \quad \pm(1, 0, 7), \quad \pm(1, 2, 3), \quad \pm(1, 2, 5), \quad \pm(1, 4, 1), \quad \pm(1, 4, 7), \\ & \pm(1, 6, 3), \quad \pm(1, 6, 5), \quad \pm(3, 0, 3), \quad \pm(3, 0, 5), \quad \pm(3, 2, 1), \quad \pm(3, 2, 7), \\ & \pm(3, 4, 3), \quad \pm(3, 4, 5), \quad \pm(3, 6, 1), \quad \pm(3, 6, 7). \end{aligned}$$

Let

$$\begin{aligned} P_1 &= \{(v_1, v_2, v_3) \in R(N, 8, 7) - R_M(N, 8, 7) \mid v_1 \equiv v_3 \pmod{4}\}, \\ P_2 &= \{(v_1, v_2, v_3) \in R(N, 8, 7) - R_M(N, 8, 7) \mid v_1 + v_3 \equiv 0 \pmod{4}\}, \end{aligned}$$

and

$$\tau_1 = \frac{1}{8} \begin{pmatrix} 4 & 8 & 4 \\ -5 & 2 & 5 \\ 2 & -4 & 6 \end{pmatrix}, \quad \tau_2 = \frac{1}{8} \begin{pmatrix} 4 & 8 & -4 \\ -5 & 2 & -5 \\ -2 & 4 & 6 \end{pmatrix}.$$

Then one may easily show that it satisfies all conditions in Theorem 2.3 and the eigenvectors of τ_1 and τ_2 are $z_1 = (1, 0, 1)$ and $z_2 = (1, 0, -1)$, respectively. Therefore $(S_{8,7} \cap Q(N)) - 15\mathbb{Z}^2 \subset Q(M)$. Since $S_{40,23} \subset Q(\text{gen}(M))$, we have $S_{40,23} \subset Q(M)$. Note that if the equation $2X^2 + 5Y^2 + 40Z^2 = 40n + 23$ has an integer solution X, Y, Z , then X, Y are odd and $X \equiv \pm 2 \pmod{5}$. Hence the equation

$$5(2x + 1)^2 + 40y^2 + 2(10z - 3)^2 = 40n + 23$$

has an integer solution for any non negative integer n , which implies that the ternary sum $p_3(x) + p_4 + 2p_7(z)$ is universal over $\mathbb{Z}$.

(vi) $p_3(x) + p_5(y) + p_7(z)$. Let

$$M = \langle 5 \rangle \perp \begin{pmatrix} 12 & -6 \\ -6 & 18 \end{pmatrix}, \quad N = \langle 2, 15, 30 \rangle.$$

Note that $\text{gen}(M) = \{[M], [N]\}$. In this case, we have

$$R(N, 4, 3) - R_M(N, 4, 3) = \pm\{(0, 1, 0), (2, 1, 2)\}.$$

If we define $P_1 = R(N, 4, 3) - R_M(N, 4, 3)$ and $\tau_1 = \frac{1}{4} \begin{pmatrix} 1 & 0 & 15 \\ 0 & -4 & 0 \\ -1 & 0 & 1 \end{pmatrix}$, it satisfies all

conditions in Theorem 2.3. The eigenvector of τ_1 is $z_1 = (0, 1, 0)$ and $Q(0, 1, 0) = 15$. Therefore $(S_{4,3} \cap Q(N)) - 15\mathbb{Z}^2 \subset Q(M)$. Since $S_{120,47} \subset Q(\text{gen}(M))$, we have $S_{120,47} \subset Q(M)$. Note that if the equation

$$3(Z - 2X)^2 + 5Y^2 + 15Z^2 = 12X^2 + 5Y^2 + 18Z^2 - 12ZX = 120n + 47$$

has an integer solution X, Y, Z , then $Z - 2X \equiv Y \equiv Z \equiv 1 \pmod{2}$, $Y \not\equiv 0 \pmod{3}$ and $Z - X \equiv \pm 2 \pmod{5}$. Hence the equation

$$15(2x - 1)^2 + 5(6y - 1)^2 + 3(10z - 3)^2 = 120n + 47$$

has an integer solution for any non negative integer n , which implies that the ternary sum $p_3(x) + p_5(y) + p_7(z)$ is universal over $\mathbb{Z}$ by (2.1).

(vii) $p_3(x) + 2p_5(y) + p_7(z)$. Let

$$M = \langle 5 \rangle \perp \begin{pmatrix} 9 & 3 \\ 3 & 21 \end{pmatrix}, \quad N = \begin{pmatrix} 6 & 3 \\ 3 & 9 \end{pmatrix} \perp \langle 20 \rangle.$$

Note that $\text{gen}(M) = \{[M], [N]\}$. One may easily show by a direct computation that $N <_{15,11} M$. Since $S_{60,26} \subset Q(\text{gen}(M))$, we have $S_{60,26} \subset Q(M)$. Now consider the equation

$$\begin{aligned} 120n + 52 &= 3(Z - 4X)^2 + 10Y^2 + 15Z^2 \\ &= 48X^2 + 18Z^2 + 10Y^2 - 24XZ = f(X, Y, Z). \end{aligned}$$

Since f is isometric to $2M$, the above equation has an integer solution X, Y, Z for any non negative integer n . Furthermore, since $Z - 4X \equiv Y \equiv Z \equiv 1 \pmod{2}$, $Y \not\equiv 0 \pmod{3}$ and $Z - 4X \equiv \pm 2 \pmod{5}$, the equation

$$15(2x - 1)^2 + 10(6y - 1)^2 + 3(10z - 3)^2 = 120n + 52$$

has an integer solution for any non negative integer n . Therefore the ternary sum $p_3(x) + 2p_5(y) + p_7(z)$ is universal over $\mathbb{Z}$.

(viii) $p_3(x) + p_7(y) + 2p_7(z)$. Let

$$M = \begin{pmatrix} 3 & 0 & 1 \\ 0 & 5 & 0 \\ 1 & 0 & 17 \end{pmatrix}, \quad N = \langle 2, 5, 25 \rangle.$$

Note that $\text{gen}(M) = \{[M], [N]\}$. One may easily show by a direct computation that $N <_{8,0} M$. Since $S_{40,32} \subset Q(\text{gen}(M))$, we have $S_{40,32} \subset Q(M)$. Now consider the equation

$$\begin{aligned} 40n + 32 &= (Y - 5X)^2 + 2Y^2 + 5Z^2 \\ &= 25X^2 + 3Y^2 + 5Z^2 - 10XY = f(X, Y, Z). \end{aligned}$$

Since f is isometric to M , the above equation has an integer solution for any non negative integer n . Therefore the equation $g(X, Y, Z) = X^2 + 2Y^2 + 5Z^2 = 40n + 24$ has an integer solution X, Y, Z with $X \equiv Y \pmod{5}$. Furthermore all of X, Y and Z have same parity. Suppose that $(x, y, z) = (a, b, c)$ is an integer solution of $g(x, y, z) = 40n + 32$ such that $a \equiv b \pmod{5}$ and $a \equiv b \equiv c \equiv 0 \pmod{2}$. Then $a \equiv c \pmod{4}$ and there is another representation

$$g\left(\frac{-3a + 2b + 5c}{4}, \frac{a - 2b + 5c}{4}, \frac{-a - 2b - c}{4}\right) = g(a, b, c) = 40n + 32$$

satisfying

$$\frac{-3a + 2b + 5c}{4} \equiv \frac{a - 2b + 5c}{4} \pmod{5}.$$

Since $\tau = \frac{1}{4} \begin{pmatrix} -3 & 2 & 5 \\ 1 & -2 & 5 \\ -1 & -2 & -1 \end{pmatrix}$ has an infinite order, there is a positive integer m such that each component of the vector $\tau^m(a, b, c) = (a_m, b_m, c_m)$ is odd and $a_m \equiv b_m \pmod{5}$, unless (a, b, c) is not an eigenvector of τ . Note that $(a, b, c) = (-2t, t, 0)$ is an eigenvector of τ and $g(-2t, t, 0) = 6t^2 \not\equiv 2 \pmod{5}$. Hence for any non negative integer n , $g(X, Y, Z) = 40n + 32$ has an integer solution X, Y, Z such that $X \equiv Y \pmod{5}$ and all of X, Y and Z are odd. Therefore the equation

$$5(2x - 1)^2 + (10y - 3)^2 + 2(10z - 3)^2 = 40n + 32$$

has an integer solution for any non negative integer n , which implies that the ternary sum $p_3(x) + p_7(y) + 2p_7(z)$ is universal over $\mathbb{Z}$ by (2.1).

(ix) $p_3(x) + 2p_3(y) + 2p_8(z)$. Let

$$M = \langle 3, 6, 16 \rangle, \quad N = \langle 1 \rangle \perp \begin{pmatrix} 18 & 6 \\ 6 & 18 \end{pmatrix}.$$

Note that $\text{gen}(M) = \{[M], [N]\}$. By a direct computation, we may easily check that $R(N, 3, 1) - R_M(N, 3, 1) = \pm\{(1, 0, 0)\}$. If we define

$$P_1 = \pm\{(1, 0, 0)\} \quad \text{and} \quad \tau_1 = \frac{1}{3} \begin{pmatrix} 3 & 0 & 0 \\ 0 & -2 & -3 \\ 0 & 3 & 0 \end{pmatrix},$$

then it satisfies all conditions in Theorem 2.3. The eigenvector of τ_1 is $z_1 = (1, 0, 0)$ and $Q(1, 0, 0) = 1$. Since $S_{24,1} \subset Q(\text{gen}(M))$, we have $S_{24,1} \subset Q(M)$, unless $24n + 1$ is not a square of an integer. Since N is contained in the spinor genus of M and 1 is represented by N , every square of an integer that has a prime divisor bigger than 3 is represented by both M and N by Lemma 2.4. Therefore every integer of the form $24n + 1$ for some positive integer n is represented by M . If the equation $3X^2 + 6Y^2 + 16Z^2 = 24n + 25$ has an integer solution X, Y, Z , then X, Y are odd and Z is not divisible by 3. Hence the equation

$$3(2x - 1)^2 + 6(2y - 1)^2 + 16(3z - 1)^2 = 24n + 25$$

has an integer solution for any non negative integer n , which implies that the ternary sum $p_3(x) + 2p_3(y) + 2p_8(z)$ is universal over $\mathbb{Z}$ by (2.1).

(x) $p_3(x) + p_7(y) + p_8(z)$. Let

$$M = \begin{pmatrix} 6 & 3 \\ 3 & 9 \end{pmatrix} \perp \langle 20 \rangle, \quad N = \langle 5 \rangle \perp \begin{pmatrix} 9 & 3 \\ 3 & 21 \end{pmatrix}.$$

Note that $\text{gen}(M) = \{[M], [N]\}$. One may show by a direct computation that $N \prec_{30,11} M$. Since $S_{60,41} \subset Q(\text{gen}(M))$, we have $S_{60,41} \subset Q(M)$. Hence the equation

$$120n + 82 = 3(Y - 2X)^2 + 15Y^2 + 40Z^2 = 12X^2 + 18Y^2 + 40Z^2 - 12XY$$

has an integer solution X, Y, Z for any non negative integer n . Since $Y - 2X \equiv Y \equiv 1 \pmod{2}$, $Y - 2X \equiv \pm 2 \pmod{5}$ and $Z \not\equiv 0 \pmod{3}$, the equation

$$15(2x - 1)^2 + 3(10y - 3)^2 + 40(3z - 1)^2 = 120n + 82$$

has an integer solution for any non negative integer n . Therefore by (2.1), the ternary sum $p_3(x) + p_7(y) + p_8(z)$ is universal over $\mathbb{Z}$.

(xi) $p_3(x) + 2p_3(y) + p_9(z)$. Let

$$M = \langle 1 \rangle \perp \begin{pmatrix} 21 & 14 \\ 14 & 28 \end{pmatrix}, \quad N = \langle 1, 7, 14 \rangle, \quad N' = \langle 2, 7, 7 \rangle.$$

Note that $\text{gen}(N) = \{[N], [N']\}$. One may easily show by a direct computation that $N' <_{8,6} N$. Since $S_{56,46} \subset Q(\text{gen}(N))$, we have $S_{56,46} \subset Q(N)$. Now, one may show by a direct computation that

$$R(N, 8, 6) - R_M(N, 8, 6) = \pm\{(0, 0, 1), (0, 0, 3), (4, 4, 1), (4, 4, 3)\}.$$

If we define $P_1 = R(N, 8, 6) - R_M(N, 8, 6)$ and $\tau_1 = \frac{1}{8} \begin{pmatrix} 1 & 21 & 0 \\ -3 & 1 & 0 \\ 0 & 0 & -8 \end{pmatrix}$, then it

satisfies all conditions in Theorem 2.3. The eigenvector of τ_1 is $z_1 = (0, 0, 1)$ and $Q(0, 0, 1) = 14$. Since $(S_{8,6} \cap Q(N)) - 14\mathbb{Z}^2 \subset Q(M)$ by Theorem 2.3, we have $S_{56,46} \subset Q(M)$. Hence the equation

$$56n + 46 = X^2 + 7(Z - 2Y)^2 + 14Z^2 = X^2 + 28Y^2 + 21Z^2 - 28YZ$$

has an integer solution X, Y, Z for any non negative integer n . Since $X \equiv Z - 2Y \equiv Z \equiv 1 \pmod{2}$ and $X \equiv \pm 2 \pmod{7}$, the equation

$$7(2x - 1)^2 + 14(2y - 1)^2 + (14z - 5)^2 = 56n + 46$$

has an integer solution for any non negative integer n , which implies that the ternary sum $p_3(x) + 2p_3(y) + p_9(z)$ is universal over $\mathbb{Z}$ by (2.1).

(xii) $p_3(x) + p_4(y) + p_9(z)$. Let

$$M = \begin{pmatrix} 2 & 1 \\ 1 & 4 \end{pmatrix} \perp \langle 28 \rangle, \quad N = \begin{pmatrix} 4 & 2 & 2 \\ 2 & 8 & 1 \\ 2 & 1 & 8 \end{pmatrix}.$$

Note that $\text{gen}(M) = \{[M], [N]\}$. One may easily show by a direct computation that $N <_{4,0} M$. Since $S_{28,16} \subset Q(\text{gen}(M))$, we have $S_{28,16} \subset Q(M)$. Hence the equation

$$56n + 32 = (Y - 2X)^2 + 7Y^2 + 56Z^2 = 4X^2 + 8Y^2 + 56Z^2 - 4XY$$

has an integer solution for any non negative integer n . Therefore the equation $g(X, Y, Z) = X^2 + 7Y^2 + 56Z^2 = 56n + 32$ has an integer solution (a, b, c) with

$a \equiv b \pmod{2}$. Suppose that (a, b, c) is an integer solution of $g(x, y, z) = 56n + 32$ satisfying $a \equiv b \equiv 0 \pmod{2}$. Note that

$$g\left(\frac{a-2b}{2}, \frac{a+2b}{2}, c\right) = g(a, b, c) = 56n + 32.$$

Since $\tau = \frac{1}{2} \begin{pmatrix} 1 & -2 & 0 \\ 1 & 2 & 0 \\ 0 & 0 & 2 \end{pmatrix}$ has an infinite order, there is a positive integer m such

that $a_m \equiv b_m \equiv 1 \pmod{2}$, where $\tau^m(a, b, c) = (a_m, b_m, c_m)$. Since $(0, 0, t)$ is an eigenvector of τ and $g(0, 0, t) = 56t^2 \equiv 0 \pmod{7}$, the equation $g(X, Y, Z) = 56n + 32$ has an integer solution X, Y, Z such that $X \equiv Y \equiv 1 \pmod{2}$ and $X \equiv \pm 2 \pmod{7}$, for any non negative integer n . Hence the equation

$$7(2x - 1) + 56y^2 + (14z - 5)^2 = 56n + 32$$

has an integer solution for any non negative integer n . Therefore by (2.1), the ternary sum $p_3(x) + p_4(y) + p_9(z)$ is universal over $\mathbb{Z}$.

(xiii) $p_3(x) + p_5(y) + 2p_9(z)$. This case is one of the most complicate one. Let

$$M = \begin{pmatrix} 13 & 1 \\ 1 & 13 \end{pmatrix} \perp \langle 21 \rangle, \quad N = \begin{pmatrix} 10 & 5 & 2 \\ 5 & 13 & 1 \\ 2 & 1 & 34 \end{pmatrix} \quad N' = \begin{pmatrix} 10 & 4 & -3 \\ 4 & 10 & 3 \\ -3 & 3 & 45 \end{pmatrix}.$$

Note that $\text{gen}(M) = \{[M], [N], [N']\}$ and $S_{21,10} \subset Q(\text{gen}(M))$. One may easily show that $N' <_{21,10} N$. The set $R(N, 21, 10) - R_M(N, 21, 10)$ consists of 48 vectors, which is the union of the following sets:

$$\begin{aligned} P_1 &= \pm\{v : v \equiv (4, 0, 17), (10, 0, 11) \pmod{21}\}, \\ P_2 &= \pm\{v : v \equiv (2, 17, 19), (5, 11, 16) \pmod{21}\}, \\ P_3 &= \pm\{v : v \equiv (2, 11, 13), (5, 17, 1) \pmod{21}\}, \\ P_4 &= \pm\{v : v \equiv (1, 0, 0), (1, 12, 9), (8, 0, 0), (8, 12, 9) \pmod{21}\}, \\ \tilde{P}_1 &= \pm\{v : v \equiv (2, 9, 1), (2, 17, 2), (4, 12, 8), (5, 11, 5), \\ &\quad (5, 12, 13), (10, 9, 20) \pmod{21}\}, \\ \tilde{P}_2 &= \pm\{v : v \equiv (1, 0, 18), (1, 4, 1), (1, 12, 12), (2, 0, 13), \\ &\quad (5, 0, 1), (8, 11, 8), (8, 12, 12), (13, 0, 3) \pmod{21}\}. \end{aligned}$$

We also define

$$\tau_1 = \frac{1}{21} \begin{pmatrix} -14 & -14 & -35 \\ 14 & -7 & 14 \\ -7 & -7 & 14 \end{pmatrix}, \tau_2 = \frac{1}{21} \begin{pmatrix} -21 & -7 & 14 \\ 6 & -17 & -2 \\ 6 & 4 & 19 \end{pmatrix}, \tau_3 = \frac{1}{21} \begin{pmatrix} -15 & -1 & -34 \\ -6 & 1 & 34 \\ 6 & 13 & 1 \end{pmatrix},$$

and

$$\tau_4 = \frac{1}{21} \begin{pmatrix} -21 & -21 & 0 \\ 0 & 19 & -16 \\ 0 & 5 & 19 \end{pmatrix}, \quad \tau_5 = \frac{1}{21} \begin{pmatrix} -21 & -21 & 0 \\ 6 & 15 & 30 \\ 6 & -6 & 9 \end{pmatrix}.$$

Note that each τ_i satisfies the condition (i),(ii) of Theorem 2.3. One may easily show that P_1 and τ_1 satisfy the condition (iii) of Theorem 2.3. For any $v \in P_2$, we may easily check that $\tau_2(v) \in N$ and if $\tau_2(v) \pmod{21} \notin R_M(N, 21, 10)$, then

$$\tau_1\tau_2(v) \in N \quad \text{and} \quad \tau_1\tau_2(v) \pmod{21} \in R_M(N, 21, 10) \cup P_2.$$

For example, assume that $v = (v_1, v_2, v_3) \equiv (2, 17, 19) \pmod{21}$. Then $\tau_2(v)$ is either a good vector or congruent to $(19, 4, 19)$ modulo 21. If the latter holds, then $\tau_1\tau_2(v)$ is either a good vector or congruent to $(2, 17, 19)$ modulo 21. Note that the order of $\tau_1\tau_2$ is infinite.

For any $v' \in P_3$, one may easily check that $\tau_3(v') \in N$ and if $\tau_3(v') \pmod{21} \notin R_M(N, 21, 10) \cup P_2$, then

$$\tau_1\tau_3(v') \in N \quad \text{and} \quad \tau_1\tau_3(v') \pmod{21} \in R_M(N, 21, 10) \cup P_3.$$

Note that the order of $\tau_1\tau_3$ is infinite. Furthermore, for any $v'' \in P_4$, one may easily check that

$$\tau_4(v'') \in N \quad \text{and} \quad \tau_4(v'') \pmod{21} \in R_M(N, 21, 10) \cup P_4 \cup \tilde{P}_2.$$

Note that for any $\tilde{v}_1 \in \tilde{P}_1$,

$$\tau_1(\tilde{v}_1) \in N \quad \text{and} \quad \tau_1(\tilde{v}_1) \pmod{21} \in R_M(N, 21, 10) \cup P_2.$$

If $\tilde{v}_2 \in \tilde{P}_2$, then one of vectors in

$$\{\tau_1(\tilde{v}_2), \tau_5(\tilde{v}_2), \tau_1\tau_5(\tilde{v}_2), \tau_1^2(\tilde{v}_2), \tau_1^2\tau_5(\tilde{v}_2)\}$$

is contained both in N and in $R_M(N, 21, 10) \cup P_3$ modulo 21. Therefore, though the situation is slightly different from the theorem, we may still apply Theorem 2.3 to this case.

The primitive eigenvectors of $\tau_1, \tau_1\tau_2, \tau_1\tau_3$ and τ_4 are $(-1, 0, 1), (-1, 2, 1), (5, -4, 1)$ and $(1, 0, 0)$, respectively. Note that

$$Q(-1, 0, 1) = 40, \quad Q(-1, 2, 1) = 76, \quad Q(5, -4, 1) = 304 \quad \text{and} \quad Q(1, 0, 0) = 10.$$

Therefore if $168n + 10$ is not of the form $10t^2$ for some positive integer t , then $168n + 10$ is represented by M . Assume that $168n + 10 = 10t^2$ for some positive integer $t > 1$. Then t has a prime divisor relatively prime to $2 \cdot 3 \cdot 7$. Since $N, N' \in \text{spn}(M)$, and both N and N' represent 10, M represents $10t^2$ by Lemma 2.4. Therefore every integer of the form $168n + 10$ for a positive integer n is represented by M . Hence the equation

$$168n + 178 = 6X^2 + 7(2Y - X)^2 + 21Z^2 = 13X^2 + 28Y^2 + 21Z^2 - 28XY$$

has an integer solution X, Y, Z for any non negative integer n . Since $X \equiv 2Y - X \equiv Z \equiv 1 \pmod{2}$, $2Y - X \not\equiv 0 \pmod{3}$ and $X \equiv \pm 2 \pmod{7}$, the equation

$$21(2x - 1)^2 + 7(6y - 1)^2 + 6(14z - 5)^2 = 168n + 178$$

has an integer solution for any non negative integer n , which implies that the ternary sum $p_3(x) + p_5(y) + 2p_9(z)$ is universal over $\mathbb{Z}$.

(xiv) $2p_3(x) + p_5(y) + p_9(z)$. Let

$$M = \begin{pmatrix} 5 & 1 \\ 1 & 17 \end{pmatrix} \perp \langle 21 \rangle, \quad N = \begin{pmatrix} 12 & 0 & 6 \\ 0 & 14 & 7 \\ 6 & 7 & 17 \end{pmatrix}, \quad N' = \begin{pmatrix} 5 & 2 \\ 2 & 5 \end{pmatrix} \perp \langle 84 \rangle.$$

Note that $\text{gen}(M) = \{[M], [N], [N']\}$. One may easily show by a direct computation that $N <_{21,20} M$ and $N' <_{21,20} M$. Since $S_{84,62} \subset Q(\text{gen}(M))$, we have $S_{84,62} \subset Q(M)$. Now consider the equation

$$\begin{aligned} 168n + 124 &= 3(Y - 4X)^2 + 7Y^2 + 42Z^2 \\ &= 48X^2 + 10Y^2 + 42Z^2 - 24XY = f(X, Y, Z). \end{aligned}$$

Since f is isometric to $2M$, the above equation has an integer solution X, Y, Z for any non negative integer n . Since $Y \equiv Y - 4X \pmod{4}$, $Y - 4X$, Y and Z are all odd. Furthermore, $X \equiv \pm 2 \pmod{7}$ and $Y \not\equiv 0 \pmod{3}$. Therefore the equation

$$42(2x - 1)^2 + 7(6y - 1)^2 + 3(14z - 5)^2 = 168n + 124$$

has an integer solution for any non negative integer n , which implies that the ternary sum $2p_3(x) + p_5(y) + p_9(z)$ is universal over $\mathbb{Z}$.

(xv) $p_3(x) + p_3(y) + p_{12}(z)$. Let

$$M = \langle 4, 5, 5 \rangle, \quad N = \langle 1, 5, 20 \rangle.$$

Note that $\text{gen}(M) = \{[M], [N]\}$ and $S_{20,1} \subset Q(\text{gen}(M))$. The set $R(N, 20, 1) - R_M(N, 20, 1)$ consists of following vectors:

$$\begin{aligned} &\pm(1, 0, 0), \quad \pm(1, 0, 5), \quad \pm(1, 0, 10), \quad \pm(1, 0, 15), \quad \pm(1, 10, 0), \quad \pm(1, 10, 5), \\ &\pm(1, 10, 10), \quad \pm(1, 10, 15), \quad \pm(9, 0, 0), \quad \pm(9, 0, 5), \quad \pm(9, 0, 10), \quad \pm(9, 0, 15), \\ &\pm(9, 10, 0), \quad \pm(9, 10, 5), \quad \pm(9, 10, 10), \quad \pm(9, 10, 15). \end{aligned}$$

If we define $P_1 = R(N, 20, 1) - R_M(N, 20, 1)$ and $\tau_1 = \frac{1}{20} \begin{pmatrix} 20 & 0 & 0 \\ 0 & -12 & 32 \\ 0 & -8 & -12 \end{pmatrix}$, then

it satisfies all conditions in Theorem 2.3. The eigenvector of τ_1 is $z_1 = (1, 0, 0)$ and $Q(1, 0, 0) = 1$. Therefore if $20n + 1$ is not of the form t^2 for a positive integer t , then it is represented by M . Since N is contained in the spinor genus of M and 1 is represented by N , every square of an integer that has a prime divisor relatively prime to 10 is represented by M by Lemma 2.4. Therefore every integer of the form $20n + 1$ for a positive integer n is represented by M . Now consider the equation

$$\begin{aligned} 40n + 42 &= 5(Y - 2X)^2 + 5Y^2 + 8Z^2 \\ &= 20X^2 + 10Y^2 + 8Z^2 - 20XY = f(X, Y, Z). \end{aligned}$$

Since f is isometric to $2M$, the above equation has an integer solution X, Y, Z for any non negative integer n . Since $Y - 2X \equiv Y \equiv Z \equiv 1 \pmod{2}$ and $Z \equiv \pm 2 \pmod{5}$, the equation

$$5(2x - 1)^2 + 5(2y - 1)^2 + 8(5z - 2)^2 = 40n + 42$$

has an integer solution for any non negative integer n , which implies that the ternary sum $p_3(x) + p_3(y) + p_{12}(z)$ is universal over $\mathbb{Z}$.

(xvi) $p_3(x) + 2p_3(y) + p_{12}(z)$. Let

$$M = \langle 5, 8, 10 \rangle, \quad N = \langle 2, 5, 40 \rangle.$$

Note that $\text{gen}(M) = \{[M], [N]\}$ and $S_{40,7} \subset Q(\text{gen}(M))$. One may show by a direct computation that $R(N, 8, 7) - R_M(N, 8, 7)$ consists of the following vectors:

$$\begin{aligned} & \pm(1, 1, 0), \quad \pm(1, 1, 4), \quad \pm(1, 3, 2), \quad \pm(1, 3, 6), \quad \pm(1, 5, 2), \quad \pm(1, 5, 6), \\ & \pm(1, 7, 0), \quad \pm(1, 7, 4), \quad \pm(3, 1, 2), \quad \pm(3, 1, 6), \quad \pm(3, 3, 0), \quad \pm(3, 3, 4), \\ & \pm(3, 5, 0), \quad \pm(3, 5, 4), \quad \pm(3, 7, 2), \quad \pm(3, 7, 6). \end{aligned}$$

Let

$$P_1 = \{(v_1, v_2, v_3) \in R(N, 8, 7) - R_M(N, 8, 7) \mid v_1 - v_2 + 2v_3 \equiv 0 \pmod{8}\},$$

$$P_2 = \{(v_1, v_2, v_3) \in R(N, 8, 7) - R_M(N, 8, 7) \mid v_1 + v_2 + 2v_3 \equiv 0 \pmod{8}\},$$

and

$$\tau_1 = \frac{1}{8} \begin{pmatrix} 2 & -10 & 20 \\ -4 & -4 & -8 \\ -1 & 1 & 6 \end{pmatrix}, \quad \tau_2 = \frac{1}{8} \begin{pmatrix} 2 & 10 & 20 \\ 4 & -4 & 8 \\ -1 & -1 & 6 \end{pmatrix}.$$

Then one may easily show that it satisfies all conditions in Theorem 2.3. The eigenvectors for τ_1, τ_2 are $z_1 = (1, 1, 0)$ and $z_2 = (1, -1, 0)$, respectively and $Q(1, 1, 0) = Q(1, -1, 0) = 7$. Therefore if $40n + 7$ is not of the form $7t^2$ for some positive integer t , then $40n + 7$ is represented by M .

Assume that there is a positive integer $t > 1$ such that $40n + 7 = 7t^2$. Then t has a prime divisor relatively prime to 10. Since $N \in \text{spn}(M)$ and N represents 7, M represents $7t^2$ for any $t > 1$ by Lemma 2.4. Therefore every integer of the form $40n + 7$ for a positive integer n is represented by M . Hence the equation $5X^2 + 8Y^2 + 10Z^2 = 40n + 47$ has an integer solution X, Y, Z for any non negative integer n . Since $X \equiv Z \equiv 1 \pmod{2}$ and $Y \equiv \pm 2 \pmod{5}$, the equation

$$5(2x - 1)^2 + 10(2y - 1)^2 + 8(5z - 2)^2 = 40n + 47$$

has an integer solution for any non negative integer n . Therefore the ternary sum $p_3(x) + 2p_3(y) + p_{12}(z)$ is universal over $\mathbb{Z}$ by (2.1).

(xvii) $p_3(x) + p_5(y) + p_{13}(z)$. Let

$$M = \begin{pmatrix} 12 & 0 & 6 \\ 0 & 44 & 22 \\ 6 & 22 & 47 \end{pmatrix}, \quad N = \begin{pmatrix} 15 & 6 & 3 \\ 6 & 20 & 10 \\ 3 & 10 & 71 \end{pmatrix}, \quad N' = \begin{pmatrix} 23 & 1 & 6 \\ 1 & 23 & 6 \\ 6 & 6 & 36 \end{pmatrix}.$$

Note that $\text{gen}(M) = \{[M], [N], [N']\}$ and $S_{264,23} \subset Q(\text{gen}(M))$. One may easily show by a direct computation that $N' <_{12,11} N$. Note that $R(N, 12, 11) - R_M(N, 12, 11)$ consists of following vectors:

$$\begin{aligned} & \pm(1, 11, 0), \quad \pm(2, 0, 1), \quad \pm(2, 0, 7), \quad \pm(2, 1, 11), \quad \pm(2, 7, 5), \quad \pm(3, 2, 0), \\ & \pm(3, 10, 0), \quad \pm(4, 5, 1), \quad \pm(4, 6, 5), \quad \pm(4, 6, 11), \quad \pm(4, 11, 7), \quad \pm(5, 7, 0). \end{aligned}$$

We define a partition of $R(N, 12, 11) - R_M(N, 12, 11)$ as follows:

$$\begin{aligned} P_1 &= \{(v_1, v_2, v_3) \mid v_1 + v_2 + v_3 \equiv 0 \pmod{12}\}, \\ P_2 &= \{(v_1, v_2, v_3) \mid v_1 + v_2 + v_3 \equiv 2, 4, 8, 10 \pmod{12}\}, \\ \tilde{P}_1 &= \{(v_1, v_2, v_3) \mid v_1 + v_2 + v_3 \equiv 1 \pmod{2}, \quad v_2 \equiv 2 \pmod{4}\}, \\ \tilde{P}_2 &= \{(v_1, v_2, v_3) \mid v_1 + v_2 + v_3 \equiv 1 \pmod{2}, \quad v_2 \equiv 0 \pmod{4}\}. \end{aligned}$$

We also define

$$\tau_1 = \frac{1}{12} \begin{pmatrix} -2 & -14 & -14 \\ -10 & 2 & 2 \\ 2 & 2 & -10 \end{pmatrix}, \tau_2 = \frac{1}{12} \begin{pmatrix} -8 & -14 & -6 \\ 4 & 4 & 24 \\ 4 & -2 & -6 \end{pmatrix}, \tilde{\tau}_1 = \frac{1}{12} \begin{pmatrix} -12 & -6 & -12 \\ 0 & 6 & 24 \\ 0 & -6 & 0 \end{pmatrix},$$

and $\tilde{\tau}_2 = \tilde{\tau}_1^2$. Then one may easily show that it satisfies all conditions in Theorem 2.3. The eigenvectors for τ_1, τ_2 are $z_1 = (-1, 1, 0)$, $z_2 = (-2, -1, 1)$, respectively, and $Q(z_1) = 23$, $Q(z_2) = 143$. Therefore if $264n + 23$ is not of the form $23t^2$ for a positive integer t , then $264n + 23$ is represented by M .

Assume that $264n + 23 = 23t^2$ for some positive integer t greater than 1. Then t has a prime divisor relatively prime to $2 \cdot 3 \cdot 11$. Since $N, N' \in \text{spn}(M)$ and both N and N' represent 23, M represents $23t^2$ by Lemma 2.4. Therefore every integer of the form $264n + 23$ for a positive integer n is represented by M . Hence the equation $264n + 287 = 3(Z - 2X)^2 + 11(Z - 2Y)^2 + 33Z^2 = 12X^2 + 44Y^2 + 33Z^2 - 12XZ - 44YZ$ has an integer solution X, Y, Z for any non negative integer n . Since $Z - 2X \equiv Z - 2Y \equiv Z \equiv 1 \pmod{2}$, $Z - 2X \equiv \pm 2 \pmod{11}$ and $Z - 2Y \not\equiv 0 \pmod{3}$, the equation

$$33(2x - 1)^2 + 11(6y - 1)^2 + 3(22z - 9)^2 = 264n + 287$$

has an integer solution for any non negative integer n . Therefore the ternary sum $p_3(x) + p_5(y) + p_{13}(z)$ is universal over $\mathbb{Z}$ by (2.1).

REFERENCES

- [1] J. W. Benham, and J. S. Hsia, *On spinor exceptional representations*, Nagoya Math. J. **87**(1982), 247-260.
- [2] F. Ge and Z. W. Sun, *On some universal sums of generalized polygonal numbers*, Colloq. Math. **145**(2016), 149-155.
- [3] S. Guo, H. Pan and Z. W. Sun, *Mixed sums of squares and triangular numbers* (II), Integers **7**(2007), #A56, 5pp (electronic).
- [4] Jangwon Ju, Kyoungmin Kim and Beong-Kweon Oh, *Spinor representations of ternary quadratic forms*, submitted.
- [5] Y. Kitaoka, *Arithmetic of quadratic forms*, Cambridge University Press, 1993.
- [6] B.-K. Oh, *Regular positive ternary quadratic forms*, Acta Arith. **147**(2011), 233-243.
- [7] B.-K. Oh, *Ternary universal sums of generalized pentagonal numbers*, J. Korean Math. Soc. **48**(2011), 837-847.
- [8] B.-K. Oh and Z. W. Sun, *Mixed sums of squares and triangular numbers* (III), J. Number Theory **129**(2009), 964-969.
- [9] O. T. O'Meara, *Introduction to quadratic forms*, Springer Verlag, New York, 1963.
- [10] Z. W. Sun, *Mixed sums of squares and triangular numbers*, Acta Arith. **127**(2007), 103-113.
- [11] Z. W. Sun, *On universal sums of polygonal numbers*, Sci. China Math. **58**(2015), 1367-1396.

DEPARTMENT OF MATHEMATICAL SCIENCES, SEOUL NATIONAL UNIVERSITY, SEOUL 08826, KOREA

E-mail address: `jjw@snu.ac.kr`

DEPARTMENT OF MATHEMATICAL SCIENCES AND RESEARCH INSTITUTE OF MATHEMATICS, SEOUL NATIONAL UNIVERSITY, SEOUL 08826, KOREA

E-mail address: `bkoh@snu.ac.kr`

DEPARTMENT OF MATHEMATICAL SCIENCES, SEOUL NATIONAL UNIVERSITY, SEOUL 08826, KOREA

E-mail address: `bangnam1@snu.ac.kr`